



ASH CLINICAL PRACTICE GUIDELINES
VENOUS THROMBOEMBOLISM (VTE)

FOR PATIENTS

Testing for blood clots



The information in this pamphlet is for adults, who are not pregnant and have not had a blood clot before.


If your health care provider thinks you might have a blood clot, the health care provider will perform different tests to see if you have one. The tests you receive depend on what your risk of a blood clot is, what type of blood clot they think it is, and what tests are available.

Why you might need testing for blood clots

Your health care provider may want to test for a blood clot, if you have symptoms, such as:

- chest pain
- breathing problems, such as shortness of breath
- fast heartbeat
- swelling or redness or pain in one leg

What are the different types of blood clots and conditions my doctor could think I have?



What are blood clots?

On the outside of the body, blood clots can be good. They stop the bleeding after an injury.

When a blood clot forms inside the body, it can sometimes block a blood vessel, such as a vein or artery.

If blocked, blood may not be able to flow and organs could be damaged.

When a blood clot blocks a vein deep in the body, it may cause **deep vein thrombosis (DVT)**.

DVT can occur in different veins in the body and cause pain or swelling in that area. **Upper extremity DVT** can occur in the veins in the upper part of the body – in your neck or arm veins. **Lower extremity DVT** can occur in the veins of the leg.

When a blood clot occurs in the lungs, it is called a **pulmonary embolism (PE)**. It happens when a clot in a leg vein separates and travels to the lungs. In the lungs, it becomes wedged, and can stop the blood flow. PE can be very dangerous.

What are the tests for blood clots?

D-dimer test

A **blood test** to look for a small protein piece (D-dimer) that is formed when a blood clot dissolves in your body.

A positive test means you may have a blood clot.

VQ scan

Ventilation-perfusion scanning – lung scans in which a radioactive dye is injected or gas is breathed in to see how well air and blood are able to flow through your lungs.

CTPA

Computed tomography pulmonary angiography – CT scan to get a picture of pulmonary arteries (that transport blood to the lungs).

Ultrasound

Compression or duplex ultrasound

An ultrasound makes images to look at the blood flow in the veins in your arms and legs.

There are pros and cons to each test:



Some tests have side effects. A VQ scan or CTPA can expose you to small amounts of radiation.



Some tests may not be available at your hospital.



Some tests diagnose clots better.




Some tests cost more.



Some tests are easier to have and take less time.

The health care provider will use different tests depending on where the clot may be.



Because there are pros and cons of each test, the American Society of Hematology has recommended what tests should be used and when.

The tests you have will depend on your risk of a blood clot:

Your risk of a blood clot can be low, intermediate, or high.

Your risk could be higher if you are older, had surgery recently, had a serious injury (such as breaking a bone), are taking hormones (such as birth control pills) or are male. But your health care provider will check your past health and ask you questions to decide on your risk.



If your health care provider thinks you might have a PULMONARY EMBOLISM (PE)

**If you are at
LOW RISK OF PE**

**D-dimer test is
recommended first**

If D-dimer is positive or not available, then a VQ scan or CTPA is suggested.

**If you are at
INTERMEDIATE RISK OF PE**

**D-dimer test is
recommended first**

If D-dimer is positive or not available, then a VQ scan or CTPA is suggested.

**If you are at
HIGH RISK OF PE**

CTPA is suggested first

If CTPA is not available or possible (because of an allergy or kidney disease), a VQ scan is suggested first. If the CTPA is positive, then the D-dimer test, ultrasound or VQ scan is suggested.

If you have a positive D-dimer test, your health care provider should not diagnose you with a blood clot right away and will need to do more tests. This is because about 1/2 of the people who have a positive D-dimer test may not actually have a clot, so another test is needed.

If all your tests are positive, you probably have a pulmonary embolism (PE) and may need to be treated.





If your health care provider thinks you might have a DEEP VEIN THROMBOSIS (DVT) in the veins in your leg – lower extremity DVT

LOW RISK OF DVT IN THE LOWER BODY

D-dimer test is recommended first

If the D-dimer test is positive or not available, then an ultrasound is recommended.

INTERMEDIATE OR HIGH RISK OF DVT IN THE LOWER BODY

Ultrasound is suggested (the ultrasound may be of the whole leg or the thigh)

If the whole leg ultrasound is negative, no more tests are needed.

If the ultrasound of the thigh is negative, another ultrasound is recommended if the reason for your symptoms is still not known.

If you have a positive D-dimer test, your doctor should not diagnose you with a blood clot right away and will need to do more tests.

If all your tests are positive, you probably have a DVT in your leg, and may need to be treated.



Scan for the full recommendations





If your health care provider thinks you might have DEEP VEIN THROMBOSIS (DVT) in the upper part of the body – in the neck, or arm veins – upper extremity DVT

LOW RISK OF DVT IN THE UPPER BODY

INTERMEDIATE OR HIGH RISK OF DVT IN THE UPPER BODY

D-dimer test is recommended first

D-dimer test or ultrasound is suggested first

If the D-dimer test is positive or not available, then an ultrasound is suggested.

If the D-dimer test is positive or not available, then an ultrasound is suggested.

If you have a positive D-dimer test, your health care provider should not diagnose you with a blood clot right away and will need to do more tests.

If all your tests are positive, you probably have a DVT in the upper part of your body, and may need to be treated.



Speak with your health care provider

To understand your tests for a blood clot, you can ask about:

- the type of blood clot your health care provider thinks you might have
- whether you are at low, intermediate or high risk of the blood clot
- the tests you may have to take
- the side effects of the tests
- what will happen if you have a positive or negative test





Scan for the full
recommendations

Wendy Lim, Grégoire Le Gal, Shannon M. Bates, Marc Righini, Linda B. Haramati, Eddy Lang, Jeffrey A. Kline, Sonja Chasteen, Marcia Snyder, Payal Patel, Meha Bhatt, Parth Patel, Cody Braun, Housne Begum, Wojtek Wiercioch, Holger J. Schünemann, Reem A. Mustafa; American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Adv* 2018; 2 (22): 3226–3256.

doi: <https://doi.org/10.1182/bloodadvances.2018024828>

ASH guidelines are reviewed annually by expert work groups convened by ASH. Resources derived from guidelines that require updating are removed from the ASH website.

Funding and Conflicts of Interest:

Funding for this patient version and the original version of the guideline was provided by the American Society of Hematology. The contributors to this version had no conflicts of interest.



American Society of Hematology